

## Ionic Compounds, Born-Haber Cycle, and Lattice Energy

1. Draw a Born-Haber Cycle for the formation of  $\text{BaCl}_2$ . Calculate the net energy change:  $\text{Ba (s)} + \text{Cl}_2 \text{ (g)} \rightarrow \text{BaCl}_2 \text{ (s)}$

Some useful information: (All of this information can be found in the textbook)

$$E_{\text{sublimation}} = 180 \text{ kJ/mol}$$

$$\text{Bond Dissociation Energy, } D, (\text{Cl}_2 \text{ (g)}) = 242.6 \text{ kJ/mol}$$

$$E_{i1} = 502.8 \text{ kJ/mol}$$

$$E_{i2} = 965.1 \text{ kJ/mol}$$

$$E_{ea} = -349 \text{ kJ/mol}$$

$$\text{Formation of } \text{BaCl}_2; \text{Ba}^{2+} \text{ (g)} + 2 \text{Cl}^- \text{ (g)} \rightarrow \text{BaCl}_2 \text{ (s)} \quad -858.6 \text{ kJ/mol}$$

2. What is the lattice energy for  $\text{BaCl}_2$ ?
3. Which of the following has the greatest lattice energy?  
 $\text{BaCl}_2$        $\text{BaO}$        $\text{BaS}$